02450 Introduction to machine learning and data mining



DTU Compute
Institut for Matematik og Computer Science

Reading material

Tan, Steinbach and Kumar
"Introduction to Data Mining"

Section 9.2.2 + 6.1-6.3

Feedback Groups of the day:

Nicolai Hagen, Kristoffer Gamst, Sebastian Villadsen

Patrick Krolmark, Hui Jin, Mads Kjeldse

Nikolaj Nøkkentved Larsen, Martin Bøje Rasmussen

Kasper Wandahl Fogh, Devran Kücükyildiz

Malene Eppler Bak, Julie Klokkerholm, Pernille Petersen

If possible, please (1) stay to give me feedback after the second lecture today (ca. 15:00) and (2) send an email or contact me at the exercises next week with feedback/suggestions on the exercises for today.

Lecture schedule

1. Introduction
   (Tan 1.1-1.4)

Data: Feature extraction and visualization

2. Data and feature extraction
   (Tan 2.1-2.3 + B1 (+ A)
3. Measures of similarity and summary statistics
   (Tan 2.4 + 3.1-3.2 + C1-C2)
4. Data visualization
   (Tan 3.3)

Supervised learning: Classification and regression

5. Decision trees and linear regression
   (Tan 4.1-4.3 + D)
6. Overfitting and performance evaluation
   (Tan 4.4-4.6)
7. Nearest neighbor, naive Bayes, and artificial neural networks
   (Tan 5.2-5.4)
8. Ensemble methods and multi class classifiers
   (Tan 5.6-5.8)

Unsupervised learning: Clustering and density est.

9. K-means and hierarchical clustering
   (Tan 8.1-8.3+8.5.7)
10. **Mixture models and association mining**
    (Tan 9.2.2 + 6.1-6.3)
11. Density estimation and anomaly detection
    (Tan 10.1-10.4)

Machine learning and data modelling in practice

12. Putting it all together: Summary and overview
13. Mini project

Data modeling framework



After today you should be able to:

Explain the role of the parameters in the Gaussian Mixture Model (GMM) and how the parameters are updated using the EM-algorithm

Explain why cross-validation can be used for GMM

Understand and apply kernel density, K-nearest neighbour density and average relative density estimation for outlier/anomaly detection.

Imagine you observe the world for the first time!



http://www.clipartlord.com/category/baby-clip-art/











We humans are skilled at dividing objects into groups (clustering), but how do we make computers do the same?

http://commons.wikimedia.org/wiki/File:Abessinier_sorrel.jpg
http://commons.wikimedia.org/wiki/File:Cat_Eyes.jpg
http://commons.wikimedia.org/wiki/File:Black_white_cat_on_fence.jpg
http://commons.wikimedia.org/wiki/File:Golden_Retriever_Dukedestiny01.jpg
http://commons.wikimedia.org/wiki/File:MasPiri-Astro-SVE.jpg
http://commons.wikimedia.org/wiki/File:GermanShorthPtr_wb.jpg
http://commons.wikimedia.org/wiki/File:Cat002.jpg
https://commons.wikimedia.org/wiki/Dog#/media/File:BluetickCoonhound.jpg
http://commons.wikimedia.org/wiki/File:Saurier2.jpg



Group exercise

- What are the clusters below and what characterize each cluster?
- Is **k-means** well suited for modeling the clusters below?
  - Will it always find the optimum solution?
  - Can it model the sizes of the clusters?
  - Can it model the shape of the clusters?
  - How can we determine the number of clusters?





Group exercise

- What are the clusters below and what characterize each cluster?
- Is **k-means** well suited for modeling the clusters below?
  - Will it always find the optimum solution?
  - Can it model the sizes of the clusters?
  - Can it model the shape of the clusters?
  - How can we determine the number of clusters?



Normal distribution

- Probability density function describes the relative chance of a given value to occur

- Normal distribution characterized by
  - Mean
  - Variance

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$$



Multivariate Normal distribution

$$\mathcal{N}(\boldsymbol{x}|\boldsymbol{\mu},\boldsymbol{\Sigma}) = \frac{1}{\sqrt{(2\pi)^d|\boldsymbol{\Sigma}|}} e^{-\frac{1}{2}(\boldsymbol{x}-\boldsymbol{\mu})^T\boldsymbol{\Sigma}^{-1}(\boldsymbol{x}-\boldsymbol{\mu})}$$





Multivariate Normal distribution

$$\mathcal{N}(\boldsymbol{x}|\boldsymbol{\mu},\boldsymbol{\Sigma}) = \frac{1}{\sqrt{(2\pi)^d|\boldsymbol{\Sigma}|}} e^{-\frac{1}{2}(\boldsymbol{x}-\boldsymbol{\mu})^T\boldsymbol{\Sigma}^{-1}(\boldsymbol{x}-\boldsymbol{\mu})}$$

- Example: 2-dimensional Normal distribution

$$\mu = \begin{bmatrix} \bar{x}_1 \\ \bar{x}_2 \end{bmatrix}$$

$$\Sigma = \begin{bmatrix} \mathrm{var}(x_1) & \mathrm{cov}(x_1, x_2) \\ \mathrm{cov}(x_2, x_1) & \mathrm{var}(x_2) \end{bmatrix}$$



The Gaussian Mixture Model (GMM)



- Different locations $\boldsymbol{\mu}_{(k)}$
- Different shape $\boldsymbol{\Sigma}_{(k)}$
- Different sizes w_k

Data density

Sum of cluster specific densities assumed normal distributed

$$p(\mathbf{x}) = \sum_{k=1}^{K} w_k \mathcal{N}(\mathbf{x}|\mu_{(k)}, \boldsymbol{\Sigma}_{(k)})$$

$$\text{s.t.} \sum_{k=1}^{K} w_k = 1, \; w_k \geq 0$$

GMM example

$$p(\boldsymbol{x}) = 0.5\mathcal{N}(\boldsymbol{x}|\begin{bmatrix}0\\0\end{bmatrix},\begin{bmatrix}1 & 0.5\\0.5 & 1\end{bmatrix}) + 0.2\mathcal{N}(\boldsymbol{x}|\begin{bmatrix}2\\2\end{bmatrix},\begin{bmatrix}1 & -0.7\\-0.7 & 1\end{bmatrix}) + 0.3\mathcal{N}(\boldsymbol{x}|\begin{bmatrix}4\\3\end{bmatrix},\begin{bmatrix}0.2 & 0.1\\0.1 & 0.5\end{bmatrix})$$







$\boldsymbol{\mu}_{(k)}$: Cluster center (prototypical example in cluster)

$\boldsymbol{\Sigma}_{(k)}$: Shape of the cluster

w_k : Relative size/density of the cluster



Group exercise

- Consider the Gaussian mixture model (GMM)

$$p(\mathbf{x}) = \sum_{k=1}^{K} w_k \mathcal{N}(\mathbf{x}|\, \mu_{(k)}, \mathbf{\Sigma}_{(k)}) \quad \text{s.t.} \sum_{k=1}^{K} w_k = 1,\ w_k \geq 0$$

- What is the value of the integral?

$$\int p(\mathbf{x}) \mathrm{d}\mathbf{x}$$

Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change

E-step

$$p(\boldsymbol{z}_n = k|\boldsymbol{x}_n) = \frac{w_k \mathcal{N}(\boldsymbol{x}|\boldsymbol{\mu}_{(k)}, \boldsymbol{\Sigma}_{(k)})}{\sum_{K=1}^{K} w_k \mathcal{N}(\boldsymbol{x}|\boldsymbol{\mu}_{(k)}, \boldsymbol{\Sigma}_{(k)})}$$

M-step

$$N_k = \sum_{n=1}^{N} p(\boldsymbol{z}_n = k|\boldsymbol{x}_n)$$

$$\boldsymbol{\mu}_{(k)} = \frac{1}{N_k} \sum_{n=1}^{N} \boldsymbol{x}_n p(\boldsymbol{z}_n = k|\boldsymbol{x}_n)$$

$$w_k = \frac{N_k}{N}$$

$$\boldsymbol{\Sigma}_{(k)} = \frac{1}{N_k} \sum_{n=1}^{N} (\boldsymbol{x}_n - \boldsymbol{\mu}_{(k)})(\boldsymbol{x}_n - \boldsymbol{\mu}_{(k)})^\top p(\boldsymbol{z}_n = k|\boldsymbol{x}_n)$$

Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change



Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change



Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change



Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change



Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change



Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change



Gaussian mixture models, EM algorithm

Select an initial set of model parameters (mean and covariance for each cluster)

Repeat

- **Expectation**
  - For each object, calculate the probability of belonging to each distribution
- **Maximization**
  - For each probability distribution, estimate parameters by maximum likelihood

Until the parameters do not change





Group exercise

- Consider the data to the right with 16 observations.
  - What would ideally happen if we used a GMM with K=16 clusters to model the data?

- Imagine we have two **test observations** denoted $\mathbf{x}_1$ and $\mathbf{x}_2$ (red points) that are not used for training.
  - What happens to $p(\mathbf{x}_1)$ and $p(\mathbf{x}_2)$ if we use K=3 and K=16 clusters?



Mixture models

- Selecting complexity using crossvalidation



Mixture models

- Selecting complexity using crossvalidation

EM 1st iteration



Mixture models

- Selecting complexity using crossvalidation

EM 2nd iteration



Mixture models

- Selecting complexity using crossvalidation

EM 3rd iteration



Mixture models

- Selecting complexity using crossvalidation

EM 4th iteration



Mixture models

- Selecting complexity using crossvalidation

EM 5th iteration



Mixture models

- Selecting complexity using crossvalidation



Mixture models

- Selecting complexity using crossvalidation



K-means versus GMM

K-means

- No guarantee of optimal solution
- Does not model shape of clusters
- Does not model the size of clusters
- Difficult to assess the number of clusters to use particularly when there is no ground truth

Gaussian mixture model (GMM)

- No guarantee of optimal solution (even more local minima issues due to the additional model parameters)
- Models shape of cluster as ellipsoid
- Models the size of clusters
- Possible to estimate the number of components by cross-validation

GMM on Iris data using 1,2,3 and 4 components









Recap of GMM on Iris data

GMM 10 fold cross-validation on Iris data repeated five times where the five runs are plotted using box-plots.



Anomaly detection: Definition

- Given a collection of data objects
  - Each object has associated a number of features
- Detect which objects **deviate from normal** behaviour

Anomaly detection: Example

- Credit card **fraud detection**
  - Recognize dubious credit card transactions based on the transaction history of the card holder
- Network **intrusion detection**
  - Detect hacker attacks, web crawlers etc.
- **Ecosystem disturbances**
  - Detect hurricanes, floods droughts, heat waves and fires
- **Health and medicine monitoring**
  - Detect abnormal behaviour in populations and patients
- **Fault detection in industry systems**
  - Detect when a wind turbine performs poorly due to ice coating on blades
- Detection of **outliers** in data measurements
  - Remove erroneous measurements due to misreading from an instrument



Group exercise

- Come up with **your own definition** of an outlier / anomaly
- How can we detect outliers using some of the methods you have already learned in the course?

Data example I: Cats, Dogs and Dinosaurs



Data example II: Whisky

- 86 types of Scotch whisky
- Human ratings 1-5
- 12 taste categories
  - body, sweetness, smoky, medicinal, tobacco, honey, spicy, winey, nutty, malty, fruity, floral



PCA plot


Glenfiddich
Auchento
Clynelish
Caol Ila
Ardbeg
Talisker
Lagavulin
Dalmore
RoyalLochnagar
Mortlach
Balmenach
Aberlour
Macallan
PCA Component 1
PCA Component 2


GlenGarioch
Talisker
Aberlour
Ardbeg
Caol Ila
Clynelish
Lagavulin
Laphroig
Tomore
PCA Component 1
PCA Component 3


GlenGarioch
Aberlour
Talisker
Benriach
Mortlach
Macallan
Glendronach
Balmenach
Speyside
Tomore
PCA Component 2
PCA Component 3

Dendrogram

- Dendrograms can be used to visualize relative distances between the observations

Data I: Cats, Dogs and Dinosaurs



Data II: Whisky



Density based techniques: Univariate normal distribution

- Map attribute to standard Normal variable

$$z = \frac{x - \mu}{\sigma}$$

- Choose a threshold

c	p(|z|>c)
1.0	0.3173
1.5	0.1336
2.0	0.0455
2.5	0.0124
3.0	0.0027
3.5	0.0005
4.0	0.0001



Normal distribution

- Map attribute to standard Normal variable

$$z = \frac{x - \mu}{\sigma}$$

- Choose a threshold

c	p(|z|>c)
1.0	0.3173
1.5	0.1336
2.0	0.0455
2.5	0.0124
3.0	0.0027
3.5	0.0005
4.0	0.0001



$$\mu = 1.55$$
$$\sigma = 0.99$$

Normal distribution

- Map attribute to standard Normal variable

$$z = \frac{x - \mu}{\sigma}$$

- Choose a threshold

c	p(|z|>c)
1.0	0.3173
1.5	0.1336
2.0	0.0455
2.5	0.0124
3.0	0.0027
3.5	0.0005
4.0	0.0001



Normal distribution

- Map attribute to standard Normal variable

- Choose a threshold

$$z = \frac{x - \mu}{\sigma}$$

$$p(|z| > c) = 0.001$$

$$c = 3.2905$$

Data I: Cats, Dogs and Dinosaurs



Data II: Whisky



Note: Assumes attributes follow a normal distribution which may not be a valid assumption!

Normal distribution

- Map attribute to standard Normal variable

$$z = \frac{x - \mu}{\sigma}$$

- Choose a threshold

$$p(|z| > c) = 0.001$$
$$c = 3.2905$$



Approaches to anomaly detection

- **Density-based techniques**
  - Estimate the density of data objects
  - Outliers are:
    - Data objects in low density area

- **We can of course use the GMM to evaluate the density of test data.**
  - why not on the training data?

- **Approaches we will presently also consider**:
  - Kernel density estimation
  - Inverse average distance to K nearest neighbours (KNN density)
  - Average relative KNN density

Density based techniques: Kernel Density Estimator

Recall the Gaussian Mixture Model (GMM)

Data density

Sum of cluster specific densities assumed normal distributed

$$p(\boldsymbol{x}) = \sum_{K=1}^{K} w_k \mathcal{N}(\boldsymbol{x}|\boldsymbol{\mu}_{(k)}, \boldsymbol{\Sigma}_{(k)})$$

$$(s.t. \sum_{k=1}^{K} w_k = 1, \quad w_k \geq 0)$$



$\boldsymbol{\mu}_{(k)}$: Cluster center (prototypical example in cluster)

$\boldsymbol{\Sigma}_{(k)}$: Shape of the cluster

w_k : Relative density of the cluster

Kernel Density estimation based on Gaussian Kernel: Consider the GMM and define a Gaussian with mean $\boldsymbol{x}_n$ and co-variance $\sigma^2 I$ around each Observation.



Let all observation weight the same, i.e. $w_n = 1/N$

$$p(\mathbf{x}) = \sum_{n=1}^{N} \frac{1}{N} \mathcal{N}(\mathbf{x}|\mathbf{x}_n, \sigma^2 \mathbf{I})$$

Only free parameter σ^2!

There is nothing special about the normal distribution. For a general mixture distribution p the general form of kernel density estimator is:

$$p(\mathbf{x}) = \sum_{n=1}^{N} \frac{1}{N} p(\mathbf{x}|\mathbf{x}_n, \theta)$$

This may be useful if x is discrete or non-negative.



How do we determine σ^2?

Data I: Cats, Dogs and Dinosaurs





How do we determine σ^2

Data I: Cats, Dogs and Dinosaurs



Density of test set based on leave-one-out cross validation



Optimal $\sigma^2=1.55$

$\sigma^2=0.01$



$\sigma^2=0.1$



$\sigma^2=1$



$\sigma^2=5$



Etimated leave-one-out density evaluated at each observation





Etimated density evaluated at each observation

Data II: Whisky



Density of test set based on leave-one-out cross validation



Optimal $\sigma^2 = 0.49$

leave-one-out densities



Data II: Whisky

Balmenach
Glen Garioch
Aberlour
Tomore
Caolla
…



Inverse distance density estimation

- **Distance based measure of density**
  - Density is inverse proportional to avereage distance to k nearest neighbors
  - Density is low if nearest neighbors are far away

$$\text{density}_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K) = \frac{1}{\frac{1}{K}\sum_{\boldsymbol{x}' \in N_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)} d(\boldsymbol{x}_i, \boldsymbol{x}')}$$

- **Relative density**
  - Density compared to density at nearest neighbors

$$\text{ard}_{\boldsymbol{X}}(\boldsymbol{x}_i, K) = \frac{\text{density}_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)}{\frac{1}{K}\sum_{\boldsymbol{x}_j \in N_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)} \text{density}_{\boldsymbol{X}_{\setminus j}}(\boldsymbol{x}_j, K)}$$

$$N_{\boldsymbol{X}}(\boldsymbol{x}, K) = \{\text{The } K \text{ observations in } \boldsymbol{X} \text{ which are nearest to } \boldsymbol{x}\}$$

$$\boldsymbol{X}_{\setminus i}^T = \left[\boldsymbol{x}_1\ \boldsymbol{x}_2 \cdots \boldsymbol{x}_{i-2}\ \boldsymbol{x}_{i-1}\ \boldsymbol{x}_{i+1}\ \boldsymbol{x}_{i+2} \cdots\ \boldsymbol{x}_N\right]$$

These measures are taken from:
"Introduction to Data Mining" by Pang-Ning Tan, Michael Steinbach, Vipin Kumar



Consider the pairwise distance matrix given to the left. What is the density and average relative density of the first observation for K=2?

$d(x_i,x_j)$	x_1	x_2	x_3	x_4	x_5	x_6	x_7
x_1	0	2.5	2.4	4.0	0.8	0.6	3.3
x_2	2.5	0	0.6	1.6	2.9	3.0	1.1
x_3	2.4	0.6	0	1.9	3.0	2.7	1.0
x_4	4.0	1.6	1.9	0	4.5	4.6	3.8
x_5	0.8	2.9	3.0	4.5	0	1.1	3.9
x_6	0.6	3.0	2.7	4.6	1.1	0	3.8
x_7	3.3	1.1	1.0	3.8	3.9	3.8	0

$$\text{density}_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K) = \frac{1}{\frac{1}{K}\sum_{\boldsymbol{x}' \in N_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)} d(\boldsymbol{x}_i, \boldsymbol{x}')}$$

$$\text{ard}_{\boldsymbol{X}}(\boldsymbol{x}_i, K) = \frac{\text{density}_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)}{\frac{1}{K}\sum_{\boldsymbol{x}_j \in N_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)} \text{density}_{\boldsymbol{X}_{\setminus j}}(\boldsymbol{x}_j, K)}$$



Consider the pairwise distance matrix given to the left. What is the density and average relative density of the first observation for K=2?

$d(\mathbf{x}_i,\mathbf{x}_j)$	$\mathbf{x}_1$	$\mathbf{x}_2$	$\mathbf{x}_3$	$\mathbf{x}_4$	$\mathbf{x}_5$	$\mathbf{x}_6$	$\mathbf{x}_7$
$\mathbf{x}_1$	0	2.5	2.4	4.0	0.8	0.6	3.3
$\mathbf{x}_2$	2.5	0	0.6	1.6	2.9	3.0	1.1
$\mathbf{x}_3$	2.4	0.6	0	1.9	3.0	2.7	1.0
$\mathbf{x}_4$	4.0	1.6	1.9	0	4.5	4.6	3.8
$\mathbf{x}_5$	0.8	2.9	3.0	4.5	0	1.1	3.9
$\mathbf{x}_6$	0.6	3.0	2.7	4.6	1.1	0	3.8
$\mathbf{x}_7$	3.3	1.1	1.0	3.8	3.9	3.8	0

$$\text{density}_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K) = \frac{1}{\frac{1}{K}\sum_{\boldsymbol{x}' \in N_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)} d(\boldsymbol{x}_i, \boldsymbol{x}')}$$

$$\text{ard}_{\boldsymbol{X}}(\boldsymbol{x}_i, K) = \frac{\text{density}_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)}{\frac{1}{K}\sum_{\boldsymbol{x}_j \in N_{\boldsymbol{X}_{\setminus i}}(\boldsymbol{x}_i, K)} \text{density}_{\boldsymbol{X}_{\setminus j}}(\boldsymbol{x}_j, K)}$$

density($\mathbf{x}_1$,2)=[1/2(0.6+0.8)]$^{-1}$=2/1.4
density($\mathbf{x}_6$,2)=[1/2(0.6+1.1)]$^{-1}$=2/1.7
density($\mathbf{x}_5$,2)=[1/2(0.8+1.1)]$^{-1}$=2/1.9

Av. Rel. Density($\mathbf{x}_1$,2)=[2/1.4]/[1/2 (2/1.7 + 2/1.9)]=1.2817

Inverse distance density estimation

- KNN density (5 nearest neighbors)

Data I: Cats , dogs and dinosaurs





Inverse distance density estimation

- KNN density (5 nearest neighbors)

Balmenach
Laphroig
GlenGarioch
Aberlour
Lagevulin
…



Average Relative density

- Average Relative KNN density (5 nearest neighbors)

Data I: Cats , dogs and dinosaurs





Average relative density

- Average relative KNN density (5 nearest neighbors)



Results using different methods

- **Kernel Density Estimation**
  - Balmenach
  - Glen Garioch
  - Aberlour
  - Tomore
  - Caolla
- **KNN density**
  - Balmenach
  - Laphroig
  - Glen Garioch
  - Aberlour
  - Lagavulin
- **KNN average relative density**
  - Balmenach
  - Glen Garioch
  - Aberlour
  - Craigallechie
  - Miltonduff

Common: Balmenach, Glen Garioch, Aberlour

Example of exam questions

QI: What is the average relative density for observation 2 (i.e. $\mathbf{x}_2$) for K=2 nearest neighbours?

A:1/5
B:3/10
C:7/10
D:1

$d(\mathbf{x}_i,\mathbf{x}_j)$	$\mathbf{x}_1$	$\mathbf{x}_2$	$\mathbf{x}_3$	$\mathbf{x}_4$	$\mathbf{x}_5$
$\mathbf{x}_1$	0	2.0	0.2	0.9	0.2
$\mathbf{x}_2$	2.0	0	1.5	0.5	2.0
$\mathbf{x}_3$	0.2	1.5	0	1.2	1.4
$\mathbf{x}_4$	0.9	0.5	1.2	0	1.0
$\mathbf{x}_5$	0.2	2.0	1.4	1.0	0

$$\text{density}_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K) = \frac{1}{\frac{1}{K}\sum_{\boldsymbol{x}' \in N_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K)} d(\boldsymbol{x}_i, \boldsymbol{x}')}$$

$$\text{ard}_{\boldsymbol{X}}(\boldsymbol{x}_i, K) = \frac{\text{density}_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K)}{\frac{1}{K}\sum_{\boldsymbol{x}_j \in N_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K)} \text{density}_{\boldsymbol{X}\setminus j}(\boldsymbol{x}_j, K)}$$

Example of exam questions

QI: What is the average relative density for observation 2 (i.e. $\mathbf{x}_2$) for K=2 nearest neighbours?

A:1/5
B:3/10
C:7/10
D:1

$d(\mathbf{x}_i,\mathbf{x}_j)$	$\mathbf{x}_1$	$\mathbf{x}_2$	$\mathbf{x}_3$	$\mathbf{x}_4$	$\mathbf{x}_5$
$\mathbf{x}_1$	0	2.0	0.2	0.9	0.2
$\mathbf{x}_2$	2.0	0	1.5	0.5	2.0
$\mathbf{x}_3$	0.2	1.5	0	1.2	1.4
$\mathbf{x}_4$	0.9	0.5	1.2	0	1.0
$\mathbf{x}_5$	0.2	2.0	1.4	1.0	0

$$\text{density}_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K) = \frac{1}{\frac{1}{K}\sum_{\boldsymbol{x}' \in N_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K)} d(\boldsymbol{x}_i, \boldsymbol{x}')}$$

$$\text{ard}_{\boldsymbol{X}}(\boldsymbol{x}_i, K) = \frac{\text{density}_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K)}{\frac{1}{K}\sum_{\boldsymbol{x}_j \in N_{\boldsymbol{X}\setminus i}(\boldsymbol{x}_i, K)} \text{density}_{\boldsymbol{X}\setminus j}(\boldsymbol{x}_j, K)}$$

QI: C